

Effects of chain decays, radial flow and $U_A(1)$ restoration on the low-mass dilepton enhancement in $\sqrt{s_{NN}} = 200$ GeV Au+Au reactions

Márton Vargyas^{a,1}, Tamás Csörgő^{b,2}, Róbert Vértési^{c,b,3}

^a*Eötvös Loránd University, Budapest, Hungary*

^b*Wigner Research Centre of Physics, Budapest, Hungary*

^c*Nuclear Physics Institute ASCR, Řež, Czech Republic*

Abstract

In $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions PHENIX reported a significant enhancement in the low-mass region ($0.1 < m_{ee} < 0.7$ GeV) of the dielectron spectrum, which is still not fully understood. Several theoretical works and an indirect measurement suggest that, due to the possible restoration of the $U_A(1)$ part of the chiral symmetry in a hot and dense medium, the mass of the η' meson may substantially decrease. This work reports on a statistically acceptable description of the PHENIX low-mass dilepton enhancement using a radial flow dominated meson spectra, chain decays of long-lived resonances and an in-medium η' mass modification.

Keywords: Relativistic Heavy-ion Collision, RHIC, PHENIX, Dilepton spectrum, 200 GeV, Au+Au, Symmetry restoration, eta' enhancement

1. Motivation

In comparison to the expected hadronic decays, a significant excess has been measured in the $(0.1 - 0.7)$ GeV invariant electron-positron mass range of the PHENIX dilepton cocktail [1]. To quantify this deviation, the corresponding χ^2 and confidence level (CL) were calculated; $\chi^2/\text{NDF} = 50.3/12$, CL = 0.01% (for 0.12–1.2 GeV mass range). Preliminary results from STAR also show a substantial, although smaller effect in a different geometrical acceptance [2]. Although this low-mass dilepton enhancement is a subject of detailed theoretical investigations, according to a recent conference report [3] this enhancement is not yet fully understood.

Lattice results [4] suggest that the transition to chirally symmetric and deconfined matter is a smooth, cross-over type transition, where the chiral symmetry restoration as measured by chiral susceptibility happens at a significantly lower temperature than the confinement of quarks to hadrons. This was measured by the Polyakov loop: the continuum

extrapolated value for the transition temperature of the renormalized Polyakov loop is 25 ± 4 MeV higher than that of the chiral susceptibility. This result suggests an interesting possibility of chiral hadron dynamics in the 150–175 MeV temperature range. Chiral symmetry restoration may change various properties of hadrons, such as their mass, width, branching ratios and decays.

The PHENIX collaboration was the first to report on a measurement of direct photon spectrum in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions in Ref. [5], where the average temperature of the photon emitted region was found to be $219 \pm 19(\text{stat}) \pm 19(\text{syst})$ MeV, implying that the initial temperature was at least about 300 MeV, and the (kinetic) freeze-out temperature of the order of 140–170 MeV.

The in-medium drop of the mass of the η' meson was anticipated long time ago [6, 7, 8]. The η' meson is different from the other pseudoscalar mesons in the sense that due to the axial anomaly it has a very large mass ($m_{\eta'} = 958$ MeV) which prevents it from acting as the 9th pseudo-Goldstone boson. However, in a hot and dense hadronic medium where the chiral symmetry is partially restored, its mass can be reduced to near its quark model value.

The first indirect observation of in-medium chi-

¹vargyas@bnl.gov

²csorgo.tamas@wigner.mta.hu

³robert.vertesi@ujf.cas.cz

ral dynamics, based on Bose-Einstein correlation (BEC) measurements and on the analysis of $\eta' \rightarrow \pi^+\pi^-\eta \rightarrow \pi^+\pi^-(\pi^+\pi^-\pi^0)$ decay chain, suggested a significant and an at least 200 MeV in-medium mass drop of the η' mesons in these reactions [9, 10, 11]. These results triggered renewed interest in theoretical investigations of $U_A(1)$ symmetry restoration: the Witten-Veneziano formula has been generalized to finite temperature to check if the indirectly measured in-medium η' masses are in the theoretically acceptable range [12], lattice QCD calculations were performed to study the temperature dependence of the $U_A(1)$ symmetry restoration [13], nuclear bound states and partial restoration of the η' mass was investigated in Ref. [14], just to mention the some of the most inspiring new results. In this manuscript we cross-check this indirect observation, based on a hadronic decay channel, utilizing the dilepton (Dalitz) decay channel of the η' mesons.

We have to keep in mind that changes in other meson properties may also occur in the hot and dense hadronic medium. For instance the broadening of the η' width, which was reported in a recent measurement of the TAPS collaboration [15] in cold nuclear matter. Nuclear photoproduction calculations on the ρ meson suggest a substantial broadening which may account for part of the excess low-mass dileptons [16]. The NA60 Collaboration reported the first measurement of a ρ spectral function, based on a low-mass dilepton measurement in a similar kinematic range as the PHENIX low mass dilepton enhancement, in $E_{Lab} = 158$ AGeV indium-indium collisions at the CERN SPS [17]. The associated spectral function of the ρ meson is found to indicate significant broadening but within experimental errors vanishing in-medium ρ mass modification [17]. The same mechanism, however, is not able to fully account for the PHENIX low-mass dilepton enhancement in $\sqrt{s_{NN}} = 200$ GeV Au+Au reactions. Also note that, contrary to RHIC collisions, a previous analysis of NA44 S+Pb data in the BEC channel showed no significant η' mass drop at CERN SPS energies [18]. This indicates that the chiral dynamics is different at RHIC energies of $\sqrt{s_{NN}} = 60, 130$ and 200 GeV from the observations at the CERN SPS top energy of $\sqrt{s_{NN}} = 19.4$ GeV (see Ref. [6] for details on this point).

The method we utilize here is fully consistent with the methods we applied in Refs. [7,8] to obtain the indirect η' mass modification measurement

using Bose-Einstein correlations of pions from η' chain decays. As far as we know (also see Ref. [3]), this is the first statistically acceptable description of the low-mass enhancement in 200 GeV Au+Au collisions at RHIC, as we shall demonstrate below. However there are several possibilities for further improvements. Ideally all the in-medium effects should be considered simultaneously. In this analysis, however, we restrict ourselves in exploring the complex phenomenon of chiral symmetry restoration only to the mass modification of the η' meson, which is expected to be the most prominent effect especially at the lower temperatures of the hadronic phase.

The Dalitz-decay ($\eta' \rightarrow e^+e^-\gamma$) of the η' meson contributes to the low mass region (below ~ 1 GeV), where the observed excess is present [1].

2. The η' meson in a hot, dense medium

2.1. Effect of mass-modification

As a result of a possible chiral symmetry restoration the mass of the η' meson can significantly change, but this mass modification can not be observed directly. Since the η' meson decays in the vacuum with its original mass and it has a relatively small interaction cross-section, its lifetime is long, compared to the fireball. Thus we can search for this effect via its enhanced production, which can be described with a modified Hagedorn-formula [18, 19]:

$$f_{\eta'} = \left(\frac{m_{\eta'}^*}{m_{\eta'}} \right)^\alpha e^{-\frac{m_{\eta'}^* - m_{\eta'}}{T_{cond}}}, \quad (1)$$

where $m_{\eta'}$ is the vacuum mass-, $m_{\eta'}^*$ is the in-medium mass of the η' meson, while T_{cond} is the characteristic slope of the η' mesons within the formed medium. In order to explore the role of η' meson enhancement on the $\sqrt{s_{NN}} = 200$ GeV Au+Au collision data of PHENIX [1], extended EXODUS simulations were elaborated to generate different dilepton spectra corresponding to physically reasonable η' properties.

2.2. The shape of the η' spectrum

According to the conservation of energy, those η' mesons that can escape from the hot and dense hadronic medium, while regaining their mass, must lose momentum. Denoting the properties of the in-medium η' with an asterisk, while quantities without further notation correspond to the free η' , the




Figure 1: The reproduced PHENIX dilepton spectrum of Ref. [1] and a resonance $p_T < 2$ GeV. The bottom shows the ratio of the two, proving the dominance of the low- p_T cocktail.

conservation of energy requires η' mesons to fulfill the

$$m_{\eta'}^{*2} + p_T^{*2} = m_{\eta'}^2 + p_T^2 \quad (2)$$

relation. This condition results in two different components of the η' spectrum. Some of the η' meson's transverse momenta is large enough to let them escape from the hot and dense medium to the asymptotic vacuum by decreasing their momenta and increasing their mass. Those η' mesons, however, that have low transverse momentum⁴ in the chirally modified vacuum, cannot come on mass shell thus they are trapped in that modified vacuum. They are released and come on-shell only after the dissociation of the modified vacuum state to the asymptotic vacuum. Both components realize an effective $|0\rangle_* + |\eta'\rangle_* \rightarrow |0\rangle + |\eta'\rangle$ transition, but in a different way. In the first scenario, the trapped η' mesons follow

$$f(p_T) = \frac{1}{2\pi m_{\eta'} B^{-1}} e^{-\frac{p_T^2}{2m_{\eta'} B^{-1}}} \quad (3)$$

distribution, where B^{-1} is the slope parameter of the η' condensate. The high p_T part of the η' spectrum, however, follows a spectrum similar to other hadrons, and can be described with the same hydrodynamical models.

The final η' spectrum is the sum of Eq. 3 and the hydrodynamical spectrum, which results in a

⁴Low transverse momentum means η' mesons with $p_T^* < \sqrt{m_{\eta'}^2 - m_{\eta'}^{*2}}$ as a result of conservation of energy (Eq. 2)

double exponential spectrum as used in Refs. [9, 10] and also in the current analysis.

3. Radial flow for dilepton enhancement

Another idea which might bring us closer to understand the excess is to introduce the m_T -scaling with radial flow of the mesons, which was not included in the original PHENIX analysis [1]. Radial flow is valid in the low p_T range, where it describes the particles' spectra well, and this is also the region where the dilepton spectrum has the most contribution from. On Fig. 1 the original PHENIX result [1] and a resonance ($p_T < 2$ GeV) cut result are plotted, confirming the dominance of the low- p_T part of the spectra. Thus examining the effect of the radial flow seems to be inevitable, as it might be responsible for certain parts of the excess.

As previous examinations suggested, a hydro-motivated spectrum might give a better estimate on the low- p_T regime for all mesons [19]. The following simplified hydro spectrum was chosen:

$$E \frac{d^3 N}{dp^3} = A \left(\frac{m_T}{m} \right)^\alpha e^{-\frac{m_T - m}{T_0 + m \langle u_T \rangle^2}}. \quad (4)$$

Hydro models usually describe the primordially created particles, but not their decay products. This effect is most important in case of the π^0 meson, which can be a daughter of many other hadrons. Thus we corrected the measured pion spectrum using the Core-Halo (C-H) picture [19, 20, 21, 22]. Primordial pions (together with decay products of rapidly decaying resonances that yield thermalized pions) are in the core, the inner set of a formed medium, while decay product pions of long lived resonances form the halo, farther away from the collision point. Thus if we divide the measured π^0 spectrum with

$$\frac{1}{\sqrt{\lambda}} = \frac{N_{core} + N_{halo}}{N_{core}}, \quad (5)$$

we get an estimation for the number of core pions. An upper boundary on the corrections for neutral pions was obtained by averaging the π^+ and π^- C-H corrections [23], $\lambda_H = (\lambda_{\pi^+} + \lambda_{\pi^-})/2$, while lower boundary was $\lambda_L = \frac{\sqrt{\lambda_H}}{2}(1 + \sqrt{\lambda_H})$, based on the consideration that long lived resonances do not produce more than twice as much charged as neutral pions. The results are shown in Fig. 2. From now on, pion spectra are always C-H corrected.

The hydro formula of Eq. 4 was simultaneously fitted to available $\pi^\pm$, $K^\pm$, p and $\bar{p}$ minimum bias




Figure 2: Core-Halo correction parameter, measuring the ratio of particles in the core compared to the total number of particles (including halo). The upper and the lower boundaries of the π^0 correction band are the average of charged pion corrections and half of the average charged pion corrections respectively.

data of Ref. [24], with the same constraints, e.g. the fit ranges were 0.2–1.0 GeV/ c^2 for $\pi^\pm$ and 0.1–1.0 GeV/ c^2 for $K^\pm$, p and $\bar{p}$ in $(m_T - m)$. It is shown on Fig. 3, as a function of p_T .

The fit results are: $\langle u_T \rangle^2 = 0.21 \pm 0.01$, and $T_0 = 178 \pm 4$ MeV, while the α parameter was within errors consistent with 0, hence it was fixed to 0. The fit quality is: $\chi^2/\text{NDF} = 63/49$, CL = 8.6%.

4. Dilepton cocktail simulations

4.1. Radial flow and chain decays

Every meson was generated with the hydro spectrum of Eq. 4, except for the η' . Their normalization factors, however, could not be determined in every case, due to the lack of data of the hydro validity range ($p_T < 2$ GeV). Moreover, these inclusive spectra contain the decay products as well besides the core particles. Therefore thermal model predictions of Ref. [25] were used in this work. This is a central reasoning of this analysis: from the number of core particles we were able to estimate the enhanced production of η' mesons, and we can conclude the in-medium mass of η' from this enhancement by substituting it into Eq. 1.

This different approach requires the introduction of chain decays: all the relevant decay chains of the mesons contributing to the low-mass dilepton




Figure 3: Simultaneous fit of Eq. 4. for $\pi^\pm$, $K^\pm$, p and $\bar{p}$ data with the constraint from Ref. [24], e.g. the fit ranges are 0.2–1.0 GeV/ c^2 for $\pi^\pm$ and 0.1–1.0 GeV/ c^2 for $K^\pm$, p and $\bar{p}$ in $(m_T - m)$. Pions are Core-Halo corrected.

spectrum were taken into account. The simulation tool was EXODUS, a Monte-Carlo-based event and decay generator already fine-tuned to the PHENIX acceptance.

The simulation results are shown with dashed lines on Fig. 4. The agreement with data has been already improved, the simulated dilepton spectrum deviates less from data, the $\chi^2/\text{NDF} = 24.1/12$, the confidence level CL = 1.9%.




Figure 4: Dilepton spectra with the hydro spectrum of Eq. 4, including chain decays. The dashed lines correspond to an original η' mass scenario (e.g. to the hydro background) while the solid lines represent the best reduced mass, $m_{\eta'}^* = 520$ MeV.

4.2. Including η' mass modification

In the previous subsection, the new, hydrodynamical background was evaluated. In this one it will be combined with the exotic effect of the η' mass modification. The η' spectrum is practically determined by two parameters: the inverse slope parameter, B^{-1} (see Eq. 3) and the in-medium η' mass, $m_{\eta'}^*$. We scanned through the physically meaningful range of these two parameters in order to find the most likely setup. A simplification was introduced on this step to reduce the simulation time: we handled $m_{\eta'}^*$ as a parameter which governs only the overall size of the η' excess (see Eq. 1), while B^{-1} is responsible for the shape of the spectrum. In fact, the slope of the longer component of the excess is also influenced by the in-medium mass, but the effect was found to be small in the physically interesting in-medium mass range. Fig. 5 shows the confidence level map of these two parameters calculated for the 0.12–1.2 GeV region. The confidence level map in 5 indicates the first results of the scan of the in-medium η' mass and the slope parameter B^{-1} . This map is not conclusive for the inverse slope parameter B^{-1} in the sense that a more subtle analysis, with smaller B^{-1} bins and $m_{\eta'}^*$ spectrum-shape dependence is needed to determine an accurate value with proper systematic errors. From this preliminary scan, $B^{-1} = 55$ MeV can be obtained. In this B^{-1} bin a reasonable mass range, $480 < m_{\eta'}^* < 600$ MeV corresponds to $CL \geq 15\%$. Regardless of B^{-1} , within the observed range, $m_{\eta'}^* < 400$ MeV can be ruled out with 99.9% confidence level.

Fig. 4 also shows the best cocktail ($B^{-1} = 55$ MeV, $m_{\eta'}^* = 520$ MeV) with a solid line. The agreement is further improved to $\chi^2/NDF = 15.2/12$, $CL = 23\%$. Fig. 6 shows the dilepton cocktail for different $m_{\eta'}^*$ values.

5. Conclusions

The current analysis is aimed at the understanding of the low-mass enhancement of the PHENIX dilepton spectrum of Ref. [1]. A more detailed description can be found in Ref. [26]. As a first step, a different hadronic cocktail was evaluated based on the fact that low- p_T resonances dominate the dilepton spectrum, as it was examined and its absolute normalization detailed in Section 3.

As far as we know (see Ref. [3]), this report is the first that is able to give a statistically acceptable description of the low-mass dilepton enhancement in




Figure 5: Preliminary confidence level map of the two η' spectrum parameters; B^{-1} and $m_{\eta'}^*$. $CL > 0.01\%$ is plotted.




Figure 6: Dilepton cocktail for different in-medium η' mass values.

200 GeV Au+Au collisions at RHIC. Apparently, radial flow and chain decays have to be taken into account given that this low-mass dilepton enhancement is in the domain of soft physics with $p_T < 2$ GeV. These effects together provide a data description with $CL=1.9\%$. Adding in-medium η' mass modification increases the quality of agreement, the best values correspond to in-medium η' mass of 520 MeV, with a $CL = 23\%$. Extremely low η' masses below 400 MeV can be excluded with 99.9 % confidence level. The current results summarize the status of our research during August 2012 [27]. In this work, the error analysis and other systematic studies are still preliminary, and further work is needed to investigate in more detail the transverse momentum and centrality dependence of our findings.

Acknowledgement

This research was partially supported by the Hungarian OTKA grant NK 101438, and by the grant CZ.1.07/2.3.00/20.0207 of the ESF programme; Education for Competitiveness Operational Programme.

References

- [1] A. Adare *et al.* [PHENIX], Phys. Rev. C **81**, 034911 (2010).
- [2] J. Zhao [STAR], J. Phys. G **38**, 124134 (2011).
- [3] I. Tserruya (for the PHENIX Collaboration) plenary talk at the Quark Matter 2012 International Conference, Washington DC. <http://www.phenix.bnl.gov/phenix/WWW/talk/archive/2012/QM12/t2020.pptx>
- [4] Z. Fodor and S. D. Katz, arXiv:0908.3341 [hep-ph], in Landolt-Boernstein Volume 1-23A.
- [5] A. Adare *et al.* [PHENIX], Phys. Rev. Lett. **104** 132301 (2010).
- [6] J. I. Kapusta, D. Kharzeev and L. D. McLerran, Phys. Rev. D **53**, 5028 (1996).
- [7] Z. Huang and X. -N. Wang, Phys. Rev. D **53**, 5034 (1996).
- [8] T. Kunihiro *et al.*, Phys. Rev. B **219**, 363 (1989).
- [9] T. Csörgő, R. Vértési and J. Sziklai, Phys. Rev. Lett. **105** 182301 (2010).
- [10] R. Vértési, T. Csörgő and J. Sziklai, Phys. Rev. C **38** 054903 (2011).
- [11] R. Vértési, T. Csörgő and J. Sziklai, Nucl. Phys. A **830**, 631C (2009).
- [12] Y. Kwon, S. H. Lee, K. Morita and G. Wolf, Phys. Rev. D **86** 034014 (2012).
- [13] A. Bazavov *et al.*, [HotQCD Collaboration], arXiv:1205.3535 [hep-lat].
- [14] D. Jido, H. Nagahiro and S. Hirenzaki, Phys. Rev. C **85** (2012) 032201.
- [15] M. Nanova [CBELSA/TAPS], arXiv:1109.4029 [nucl-ex].
- [16] F. Riek, R. Rapp, Y. Oh and T. -S. H. Lee, Phys. Rev. C **82**, 015202 (2010).
- [17] R. Arnaldi *et al.* [NA60 Collaboration], Phys. Rev. Lett. **96** 162302 (2006).
- [18] S.E. Vance, T. Csörgő, D. Kharzeev, Phys. Rev. Lett. **81**, 2205 (1998).
- [19] T. Csörgő and B. Lörstad, Phys. Rev. C **54**, 1390 (1996).
- [20] R. Lednicky and M. I. Podgoretsky, Sov. J. Nucl. Phys. **30**, 432 (1979) [Yad. Fiz. **30**, 837 (1979)].
- [21] J. Bolz, U. Ornik, M. Plumer, B. R. Schlei and R. M. Weiner, Phys. Lett. B **300**, 404 (1993).
- [22] T. Csörgő, B. Lörstad and J. Zimányi, Z. Phys. C **71** (1996) 491 [hep-ph/9411307].
- [23] S. S. Adler *et al.* [PHENIX], Phys. Rev. Lett. **93**, 152302 (2004) [nucl-ex/0401003].
- [24] S. S. Adler *et al.* [PHENIX], Phys. Rev. C **69**, 034909 (2004) [nucl-ex/0307022].
- [25] M. Kaneta and N. Xu, nucl-th/0405068.
- [26] M. Vargyas. Analysis of low-mass dilepton enhancement. Lambert Academic Publishing. 2012. ISBN 978-3-659-24230-4.
- [27] M. Vargyas, T. Csörgő, R. Vértési, poster presented at the Quark Matter 2012 International Conference, <https://www.phenix.bnl.gov/phenix/WWW/talk/archive/2012/QM12/t2003.pdf>